

Name _____

Physics 110 Quiz #1, September 20, 2019

Please show all work, thoughts and/or reasoning in order to receive partial credit. The quiz is worth 10 points total.

I affirm that I have carried out my academic endeavors with full academic honesty.

Suppose that you reside on the third floor of a multi-story residence hall. One day you are looking out of your window and you notice water balloons falling past and that they strike the ground located $15m$ below $0.83s$ seconds after they pass you.

- a. With what velocity were the water balloons going when they passed by your window?

$$y_f = y_i + v_{iy}t + \frac{1}{2}a_y t^2$$

$$y_f = y_i + v_{iy}t - \frac{1}{2}gt^2$$

$$0m = 15m + v_{iy}(0.83s) - 4.9\frac{m}{s^2}(0.83s)^2$$

$$v_{iy} = -14\frac{m}{s}$$

- b. With what velocity would the water balloons impact the sidewalk below?

$$v_{fy} = v_{iy} + a_y t = -14\frac{m}{s} - 9.8\frac{m}{s^2}(0.83s) = -22.1\frac{m}{s}$$






- c. If the water balloons were dropped from rest, from what floor were the balloons being dropped? Assume that each floor is $5m$ high.

$$v_{fy}^2 = v_{iy}^2 + 2a_y \Delta y = -2g\Delta y$$

$$\Delta y = -\frac{v_{fy}^2}{2g} = -\frac{(-22.1\frac{m}{s})^2}{2 \times 9.8\frac{m}{s^2}} = -24.9m$$

$$\Delta y = y_f - y_i = -y_i \sim -25m \times \frac{1\text{floor}}{5m} = 5^{\text{th}} \text{ floor}$$

- d. Which of the following would give a possible trajectory for the water balloons as a function of time? Take up away from the ground as the positive y-axis.



- e. Suppose that for some unexplained reason that one of the water balloons didn't break when it hit the ground. You go pick it up and decide that it would be a good idea to throw it at a friend walking down the sidewalk at you. If you accelerate the balloon from rest over a distance of about 1.7m during your throw, with what speed will the water balloon leave your hand?

$$v_{fx}^2 = v_{ix}^2 + 2a_x \Delta x$$

$$v_{fx} = \sqrt{2a_x (1.7\text{m})} = \sqrt{3.4a} = 1.85\sqrt{a}$$

Physics 110 Formulas

Equations of Motion

$$\text{displacement: } \begin{cases} x_f = x_i + v_{ix}t + \frac{1}{2}a_x t^2 \\ y_f = y_i + v_{iy}t + \frac{1}{2}a_y t^2 \end{cases}$$

$$\text{velocity: } \begin{cases} v_{fx} = v_{ix} + a_x t \\ v_{fy} = v_{iy} + a_y t \end{cases}$$

$$\text{time-independent: } \begin{cases} v_{fx}^2 = v_{ix}^2 + 2a_x \Delta x \\ v_{fy}^2 = v_{iy}^2 + 2a_y \Delta y \end{cases}$$

Uniform Circular Motion

$$F_r = ma_r = m \frac{v^2}{r}; \quad a_r = \frac{v^2}{r}$$

$$v = \frac{2\pi r}{T}$$

$$F_G = G \frac{m_1 m_2}{r^2}$$

Geometry /Algebra

Circles Triangles Spheres

$$C = 2\pi r \quad A = \frac{1}{2}bh \quad A = 4\pi r^2$$

$$A = \pi r^2 \quad V = \frac{4}{3}\pi r^3$$

Quadratic equation : $ax^2 + bx + c = 0$,

$$\text{whose solutions are given by : } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Vectors

$$\text{magnitude of a vector: } v = |\vec{v}| = \sqrt{v_x^2 + v_y^2}$$

$$\text{direction of a vector: } \phi = \tan^{-1}\left(\frac{v_y}{v_x}\right)$$

Useful Constants

$$g = 9.8 \text{ m/s}^2 \quad G = 6.67 \times 10^{-11} \text{ Nm}^2/\text{kg}^2$$

$$N_A = 6.02 \times 10^{23} \text{ atoms/mole} \quad k_B = 1.38 \times 10^{-23} \text{ J/K}$$

$$\sigma = 5.67 \times 10^{-8} \text{ W/m}^2\text{K}^4 \quad v_{\text{sound}} = 343 \text{ m/s}$$

Linear Momentum/Forces

$$\vec{p} = m\vec{v}$$

$$\vec{p}_f = \vec{p}_i + \vec{F}\Delta t$$

$$\vec{F} = m\vec{a}$$

$$\vec{F}_s = -k\vec{x}$$

$$F_f = \mu F_N$$

Work/Energy

$$K_t = \frac{1}{2}mv^2$$

$$K_r = \frac{1}{2}I\omega^2$$

$$U_g = mgh$$

$$U_s = \frac{1}{2}kx^2$$

$$W_T = Fd\cos\theta = \Delta E_T$$

$$W_R = \tau\theta = \Delta E_R$$

$$W_{\text{net}} = W_R + W_T = \Delta E_R + \Delta E_T$$

$$\Delta E_R + \Delta E_T + \Delta U_g + \Delta U_s = 0$$

$$\Delta E_R + \Delta E_T + \Delta U_g + \Delta U_s = -\Delta E_{\text{diss}}$$

Heat

$$T_C = \frac{5}{9}[T_F - 32]$$

$$T_F = \frac{9}{5}T_C + 32$$

$$L_{\text{new}} = L_{\text{old}}(1 + \alpha\Delta T)$$

$$A_{\text{new}} = A_{\text{old}}(1 + 2\alpha\Delta T)$$

$$V_{\text{new}} = V_{\text{old}}(1 + \beta\Delta T) : \beta = 3\alpha$$

$$PV = Nk_B T$$

$$\frac{3}{2}k_B T = \frac{1}{2}mv^2$$

$$\Delta Q = mc\Delta T$$

$$P_C = \frac{\Delta Q}{\Delta t} = \frac{kA}{L}\Delta T$$

$$P_R = \frac{\Delta Q}{\Delta t} = \epsilon\sigma A\Delta T^4$$

$$\Delta U = \Delta Q - \Delta W$$

Rotational Motion

$$\theta_f = \theta_i + \omega_i t + \frac{1}{2}\alpha t^2$$

$$\omega_f = \omega_i + \alpha t$$

$$\omega_f^2 = \omega_i^2 + 2\alpha\Delta\theta$$

$$\tau = I\alpha = rF$$

$$L = I\omega$$

$$L_f = L_i + \tau\Delta t$$

$$\Delta s = r\Delta\theta : v = r\omega : a_t = r\alpha$$

$$a_r = r\omega^2$$

Fluids

$$\rho = \frac{M}{V}$$

$$P = \frac{F}{A}$$

$$P_d = P_0 + \rho g d$$

$$F_B = \rho g V$$

$$A_1 v_1 = A_2 v_2$$

$$\rho_1 A_1 v_1 = \rho_2 A_2 v_2$$

$$P_1 + \frac{1}{2}\rho v_1^2 + \rho g h_1 = P_2 + \frac{1}{2}\rho v_2^2 + \rho g h_2$$

Simple Harmonic Motion/Waves

$$\omega = 2\pi f = \frac{2\pi}{T}$$

$$T_s = 2\pi\sqrt{\frac{m}{k}}$$

$$T_p = 2\pi\sqrt{\frac{I}{\beta}}$$

$$v = \pm\sqrt{\frac{k}{m}}A\left(1 - \frac{x^2}{A^2}\right)^{\frac{1}{2}}$$

$$x(t) = A\sin\left(\frac{2\pi}{T}t\right)$$

$$v(t) = A\sqrt{\frac{k}{m}}\cos\left(\frac{2\pi}{T}t\right)$$

$$a(t) = -A\frac{k}{m}\sin\left(\frac{2\pi}{T}t\right)$$

$$v = f\lambda = \sqrt{\frac{F_T}{\mu}}$$

$$f_n = n f_1 = n \frac{v}{2L}$$

$$I = 2\pi^2 f^2 \rho v A^2$$

Sound

$$v = f\lambda = (331 + 0.6T) \frac{\text{m}}{\text{s}}$$

$$\beta = 10 \log \frac{I}{I_0}; \quad I_0 = 1 \times 10^{-12} \frac{\text{W}}{\text{m}^2}$$

$$f_n = n f_1 = n \frac{v}{2L}; \quad f_n = n f_1 = n \frac{v}{4L}$$