Name

Physics 111 Quiz #6, February 25, 2011

Please show all work, thoughts and/or reasoning in order to receive partial credit. The quiz is worth 10 points total.

1. Consider the arrangement of lenses shown below. Lens #1 is on the left with focal length f_1 while lens #2 is on the right with focal length f_2 . An object is placed as shown on the left of lens #1. With respect to the object, the final image will be

a. real and inverted.

- b. *real and upright*.
- c. virtual and inverted.

d. virtual and upright.



2. Consider an equilateral triangle shaped piece of glass with index of refraction ($n_{glass} = 1.5$) surrounded on all sides by air ($n_{air} = 1.0$), in which there is a beam of light incident.



a. What is the angle of refraction for the light incident on the left surface if the light is incident at 30° with respect to the normal as shown?

 $n_{air}\sin\theta_{air} = n_{glass}\sin\theta_{glass} \rightarrow 1.0\sin 30 = 1.5\sin\theta_{glass} \rightarrow \theta_{glass} = 19.5^{\circ}$

b. What is the angle (with respect to the normal) at which the light ray will emerge from the right side of the equilateral triangle? (The normal on the right side surface is drawn for you.)

From the geometry we have: $n_{glass} \sin \theta_3 = n_{air} \sin \theta_4 \rightarrow 1.5 \sin 40.5 = 1.0 \sin \theta_4 \rightarrow \theta_4 = 76.9^\circ$

Physics 111 Equation Sheet

Electric Forces, Fields and Potentials

$$\vec{F} = k \frac{Q_1 Q_2}{r^2} \hat{r}$$
$$\vec{E} = \frac{\vec{F}}{q}$$
$$\vec{E}_Q = k \frac{Q}{r^2} \hat{r}$$
$$PE = k \frac{Q_1 Q_2}{r}$$
$$V(r) = k \frac{Q}{r}$$
$$E_x = -\frac{\Delta V}{\Delta x}$$
$$W_{A,B} = q \Delta V_{A,B}$$

Magnetic Forces and Fields

 $F = qvB\sin\theta$ $F = IIB\sin\theta$ $\tau = NIAB\sin\theta = \mu B\sin\theta$ $PE = -\mu B\cos\theta$ $B = \frac{\mu_0 I}{2\pi r}$

$$\varepsilon_{induced} = -N \frac{\Delta \varphi_B}{\Delta t} = -N \frac{\Delta (DAC030)}{\Delta t}$$
Constants
 $g = 9.8 \frac{m}{s^2}$
 $le = 1.6 \times 10^{-19} C$
 $k = \frac{1}{4\pi \varepsilon_o} = 9 \times 10^9 \frac{C^2}{Nm^2}$
 $\varepsilon_o = 8.85 \times 10^{-12} \frac{Nm^2}{C^2}$
 $leV = 1.6 \times 10^{-19} J$
 $\mu_o = 4\pi \times 10^{-7} \frac{Tm}{A}$
 $c = 3 \times 10^8 \frac{m}{s}$
 $h = 6.63 \times 10^{-31} kg = \frac{0.511MeV}{c^2}$
 $m_p = 1.67 \times 10^{-27} kg = \frac{937.1MeV}{c^2}$
 $m_n = 1.69 \times 10^{-27} kg = \frac{948.3MeV}{c^2}$
 $lamu = 1.66 \times 10^{-27} kg = \frac{931.5MeV}{c^2}$
 $lamu = 1.66 \times 10^{-27} kg = \frac{931.5MeV}{c^2}$
 $Ax^2 + Bx + C = 0 \Rightarrow x = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}$

Electric Circuits

$$I = \frac{\Delta Q}{\Delta t}$$

$$V = IR = I \left(\frac{\rho L}{A}\right)$$

$$R_{series} = \sum_{i=1}^{N} R_{i}$$

$$\frac{1}{R_{parallel}} = \sum_{i=1}^{N} \frac{1}{R_{i}}$$

$$P = IV = I^{2}R = \frac{V^{2}}{R}$$

$$Q = CV = \left(\frac{\kappa \varepsilon_{0} A}{d}\right)V = (\kappa C_{0})V$$

$$PE = \frac{1}{2}QV = \frac{1}{2}CV^{2} = \frac{Q^{2}}{2C}$$

$$Q_{charge}(t) = Q_{max}\left(1 - e^{-\frac{t}{RC}}\right)$$

$$Q_{discharge}(t) = Q_{max}e^{-\frac{t}{RC}}$$

$$C_{parallel} = \sum_{i=1}^{N} C_{i}$$

$$\frac{1}{C_{vertics}} = \sum_{i=1}^{N} \frac{1}{C_{i}}$$

 $\frac{\Delta \phi_B}{\Delta \phi_B} = -N \frac{\Delta (BA \cos \theta)}{\Delta (BA \cos \theta)}$ Light as a Particle & Relativity

$$E = hf = \frac{hc}{\lambda} = pc$$

$$KE_{max} = hf - \phi = eV_{stop}$$

$$\Delta \lambda = \frac{h}{m_e c} (1 - \cos \phi)$$

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$p = \gamma mv$$

$$E_{total} = KE + E_{rest} = \gamma mc^2$$

$$E_{total}^2 = p^2 c^2 + m^2 c^4$$

$$E_{rest} = mc^2$$

$$KE = (\gamma - 1)mc^2$$

Geometry

Circles: $C = 2\pi r = \pi D$ $A = \pi r^2$ Triangles: $A = \frac{1}{2}bh$ Spheres: $A = 4\pi r^2$ $V = \frac{4}{3}\pi r^3$ Light as a Wave

$$c = f\lambda = \frac{1}{\sqrt{\varepsilon_o \mu_o}}$$

$$S(t) = \frac{energy}{time \times area} = c\varepsilon_o E^2(t) = c \frac{B^2(t)}{\mu_0}$$

$$I = S_{avg} = \frac{1}{2}c\varepsilon_o E_{max}^2 = c \frac{B_{max}^2}{2\mu_0}$$

$$P = \frac{S}{c} = \frac{Force}{Area}$$

$$S = S_o \cos^2 \theta$$

$$v = \frac{1}{\sqrt{\varepsilon\mu}} = \frac{c}{n}$$

$$\theta_{inc} = \theta_{refl}$$

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

$$\frac{1}{f} = \frac{1}{d_o} + \frac{1}{d_i}$$

$$M = \frac{h_i}{h_o} = -\frac{d_i}{d_o}$$

$$M_{total} = \prod_{i=1}^N M_i$$

$$d \sin \theta = m\lambda \text{ or } (m + \frac{1}{2})\lambda$$

$$a \sin \phi = m'\lambda$$

Nuclear Physics $E_{binding} = (Zm_p + Nm_n - m_{rest})^2$ $\frac{\Delta N}{\Delta t} = -\lambda N_o \rightarrow N(t) = N_o e^{-\lambda t}$ $A(t) = A_o e^{-\lambda t}$ $m(t) = m_o e^{-\lambda t}$ $t_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$

Misc. Physics 110 Formulae

$$\vec{F} = \frac{\Delta \vec{p}}{\Delta t} = \frac{\Delta (mv)}{\Delta t} = m\vec{a}$$
$$\vec{F} = -k\vec{y}$$
$$\vec{F}_{c} = m\frac{v^{2}}{R}\hat{r}$$
$$W = \Delta KE = \frac{1}{2}m(v_{f}^{2} - v_{i}^{2}) = -\Delta PE$$
$$PE_{gravity} = mgy$$
$$PE_{spring} = \frac{1}{2}ky^{2}$$