

Physics 111 Homework Solutions Week #2 – Friday

Tuesday, January 14, 2014

Chapter 14

Questions

- None

Multiple-Choice

- None

Problems

- None

Wednesday, January 15, 2014

Chapter 15

Questions

15.5 $\frac{V}{m} = \frac{J}{Cm} = \frac{Nm}{Cm} = \frac{N}{C}$

15.8 They are always perpendicular since if there were a component of the electric field parallel to the equipotential surface then work would be done moving a charge around the equipotential surface. This cannot be the case since the work done in moving a charge around the equipotential surface has to be zero. The only way for this to happen is to have no component of the electric field parallel to the equipotential surface, but rather only perpendicular.

15.11 Suppose that the electric field points along the positive x-axis. If the dipole is oriented so that the negative charge is to the left of the positive charge then the dipole is in a stable equilibrium, for if the charge is displaced slightly off of the axis then the dipole will experience a torque that will return to the dipole to its original configuration. However if the dipole is oriented so that the negative charge is on the right of the positive charge then the dipole is in an unstable equilibrium. Here if the dipole is displaced slightly it will experience a net torque and this torque will rotate the dipole so that the negative charge is on the left of the positive charge.

Multiple-Choice

15.2 B

15.4 A

15.5 B

15.6 A

15.7 D

Problems

15.22. Bringing each charge Q in one at a time from very far away:

i. the first charge is brought in for free since it feels no forces


ii. the second charge requires $W = -\frac{kQ}{L} = -0.9J$

iii. the next charge interacts with both of the previous and so the additional work is $W = -\frac{kQ}{L} - \frac{kQ}{\sqrt{L^2 + L^2}} = -1.54 J$, where the denominator in the second term is the diagonal distance between charges 1 and 3

iv. the last charge costs an additional $W = -\frac{kQ}{L} - \frac{kQ}{\sqrt{L^2 + L^2}} - \frac{kQ}{L} = -2.44 J$

The total work to assemble all 4 charges is then -4.88 J

15.3



- a. $V = \sum \frac{kQ}{r}$ and since equal and opposite charges are equally distant from the observation point at the origin, the two terms add up to zero – remember these are just + and – numbers, not vectors
- b. The electric fields from each charge do not cancel, but both point in the same direction (to the left) since the force from both charges on a positive test charge at the origin is to the left. Adding these up gives $E = 2\frac{kQ}{x^2}$, where $Q = 10 \mu C$ and $x = 0.1 m$, so $E = 1.8 \times 10^7 N/C$, pointing to the left.
- c. Since the potential at the origin is $V = 0$, as it is at infinity (very far away), then there is no change in V for the third charge and therefore no net work is required.

Repeating for two positive charges:

- a. $V = \frac{2kQ}{r} = 1.8 \times 10^6 V$;
- b. In this case $E = 0$ since the E fields from each charge point in opposite directions and now cancel;
- c. The net work required is $Q\Delta V = -18 J$ or 18J by an external force.

15.11 The energy is related to the charge transferred and the potential difference. We

$$\text{have } W = q\Delta V \rightarrow \Delta V = \frac{W}{q} = \frac{30 \times 10^6 J}{5C} = 6 \times 10^6 V = 6MV.$$

15.24 Rutherford Backscattering Spectrometry

- a. The electrostatic repulsion due to the two positive nuclei did work bringing the alpha particle to rest. The alpha particle loses its kinetic energy in favor of potential energy as it comes to rest at the quoted distance. Thus we have

$$\Delta KE + \Delta PE = (KE_f - KE_i) + (PE_f - PE_i) = 0 \rightarrow KE_i = PE_f$$

$$KE_i = \frac{kQ_{Au}Q_{\alpha}}{r} = \frac{9 \times 10^9 \frac{Nm^2}{C^2} \left(79e \times \frac{1.6 \times 10^{-19} C}{1e} \right) \left(2e \times \frac{1.6 \times 10^{-19} C}{1e} \right)}{9.23 \times 10^{-15} m} = 3.9 \times 10^{-12} J$$

- b. The work done accelerating the alpha particle changed its kinetic energy. Thus

$$W = \Delta KE = q\Delta V \rightarrow \Delta V = \frac{W}{q} = \frac{3.9 \times 10^{-12} J}{2e \times \frac{1.6 \times 10^{-19} C}{1e}} = 1.2 \times 10^7 V = 12 MV$$

- c. The work done is $3.9 \times 10^{-12} J$.

- d. The electrostatic repulsion due to the two positive nuclei did work bringing the alpha particle to rest. The alpha particle loses its kinetic energy in favor of potential energy as it comes to rest at the quoted distance. Thus we have

$$\Delta KE + \Delta PE = (KE_f - KE_i) + (PE_f - PE_i) = 0 \rightarrow KE_i = PE_f$$

$$KE_i = 3.3 \times 10^6 eV \times \frac{1.6 \times 10^{-19} C}{1e} = 5.28 \times 10^{-13} J$$

$$\rightarrow 5.28 \times 10^{-13} J = \frac{kQ_{Au}Q_{\alpha}}{r} = \frac{9 \times 10^9 \frac{Nm^2}{C^2} \left(79e \times \frac{1.6 \times 10^{-19} C}{1e} \right) \left(2e \times \frac{1.6 \times 10^{-19} C}{1e} \right)}{r}$$

$$\therefore r = 6.9 \times 10^{-14} m$$