

Physics 111 Homework Solutions Collected on Monday 11/3

Wednesday, October 29, 2014

Chapter 23

Questions

- none

Multiple-Choice

23.14 C

Problems

23.9 The absorption coefficient is given by

$$\ln \frac{S}{S_0} = \mu x \rightarrow \mu = \frac{1}{x} \ln \frac{S}{S_0} = \frac{1}{0.015m} \ln \frac{0.98S_0}{S_0} = -1.35m^{-1}.$$

23.10 We are given that with the same incident intensity sample 1 has a 95% transmittance, while sample 2 has an 85% transmittance. Thus for sample 1 we find that $\ln(0.95) = \mu_1 x$ and from sample 2, $\ln(0.85) = \mu_2 x$. Taking the ratio we

$$\text{find that } \frac{\mu_1}{\mu_2} = \frac{x \ln(0.95)}{x \ln(0.85)} = 0.32.$$

23.11 The CT number is given as $1000 \left[\frac{\mu - \mu_w}{\mu_w} \right] = 1000 \left[\frac{-7 \times 10^{-3} x^{-1} + 5 \times 10^{-3} x^{-1}}{-5 \times 10^{-3} x^{-1}} \right] = 400.$

μ and μ_w are obtained from

$$\mu = \frac{1}{x} \ln \frac{I}{I_0} = \frac{1}{x} \ln \frac{0.993S_0}{S_0} = -7 \times 10^{-3} x^{-1}$$

$$\mu_w = \frac{1}{x} \ln \frac{I}{I_0} = \frac{1}{x} \ln \frac{0.995S_0}{S_0} = -5 \times 10^{-3} x^{-1}.$$

Thursday, October 30, 2014

Chapters 19 & 24

Questions

24.1 Based on special relativity we know that as a particle with mass travels near the speed of light its mass increases. In order to accelerate this object from rest to a speed near that of light would require an ever increasing force (one that rapidly becomes larger by a factor of γ .) There are no known forces that could accelerate a particle with mass to the speed of light in a finite amount of time and with a finite amount of energy. So for objects with no rest mass, as they travel at the speed of light, there mass does not increase with increasing speed and we avoid these problems of accelerating the massless particles.

- 24.6 The Compton shift in wavelength for the proton and the electron are given by $\Delta\lambda_p = \frac{h}{m_p c}(1 - \cos\phi)$ and $\Delta\lambda_e = \frac{h}{m_e c}(1 - \cos\phi)$ respectively. Evaluating the ratio of the shift in wavelength for the proton to the electron, evaluated at the same detection angle ϕ , we find $\frac{\Delta\lambda_p}{\Delta\lambda_e} = \frac{m_e}{m_p} = \frac{1.67 \times 10^{-27} \text{ kg}}{9.11 \times 10^{-31} \text{ kg}} = \frac{1}{1833} = 5 \times 10^{-4}$. Therefore the shift in wavelength for the proton is smaller than the wavelength shift for the electron.

Multiple-Choice

19.16 D

19.17 A

Problems

19.15 The frequency is given as $f = \frac{c}{\lambda} = \frac{3 \times 10^8 \frac{\text{m}}{\text{s}}}{5.5 \times 10^{-7} \text{ m}} = 5.5 \times 10^{14} \text{ s}^{-1}$.

19.17 A Neodymium-YAG laser

- a. The number of photons is given as the total beam energy divided by the energy

per photon, or $\# = \frac{5J}{\frac{hc}{\lambda}} = \frac{5J}{6.63 \times 10^{-34} \text{ Js} \times 3 \times 10^8 \frac{\text{m}}{\text{s}}} \times 1.06 \times 10^{-6} \text{ m} = 1.33 \times 10^{20}$.

- b. The power in the beam is the energy delivered per unit time, or

$$P = \frac{\Delta E}{\Delta t} = \frac{5J}{1 \times 10^{-9} \text{ s}} = 5 \times 10^9 \text{ W} = 5 \text{ GW}.$$

- c. The average power output of the laser is $P = \frac{5J}{\text{pulse}} \times \frac{10 \text{ pulses}}{1 \text{ s}} = 50 \text{ W}$.

19.18 The range of frequencies is given as $f_{400\text{nm}} = \frac{c}{\lambda_{400\text{nm}}} = \frac{3 \times 10^8 \frac{\text{m}}{\text{s}}}{400 \times 10^{-9} \text{ m}} = 7.5 \times 10^{14} \text{ Hz}$

through $f_{750\text{nm}} = \frac{c}{\lambda_{750\text{nm}}} = \frac{3 \times 10^8 \frac{\text{m}}{\text{s}}}{750 \times 10^{-9} \text{ m}} = 4 \times 10^{14} \text{ Hz}$. The energies are given by

$$E = \frac{hc}{\lambda} = hf \text{ which for } 750 \text{ nm the energy is}$$

$$E = \frac{(6.63 \times 10^{-34} \text{ Js})(3 \times 10^8 \frac{\text{m}}{\text{s}})}{750 \times 10^{-9} \text{ m}} = 2.65 \times 10^{-19} \text{ J} \times \frac{1 \text{ eV}}{1.6 \times 10^{-19} \text{ J}} = 1.66 \text{ eV} \text{ while the}$$

energy for the 400 nm photon is

$$E = \frac{(6.63 \times 10^{-34} \text{ Js})(3 \times 10^8 \frac{\text{m}}{\text{s}})}{400 \times 10^{-9} \text{ m}} = 4.97 \times 10^{-19} \text{ J} \times \frac{1 \text{ eV}}{1.6 \times 10^{-19} \text{ J}} = 3.1 \text{ eV}.$$

19.20 The energy of the emitted photon is equal to the difference in energies of the two levels: $\Delta E = -1.51\text{eV} - (-3.4\text{eV}) = 1.89\text{eV} \times \frac{1.6 \times 10^{-19}\text{J}}{1\text{eV}} = 3.02 \times 10^{-19}\text{J}$. Then

setting this energy equal to $\frac{hc}{\lambda}$ we find

$$\lambda = \frac{hc}{E} = \frac{6.63 \times 10^{-34}\text{Js} \times 3 \times 10^8 \frac{\text{m}}{\text{s}}}{3.02 \times 10^{-19}\text{J}} = 6.58 \times 10^{-7}\text{m} = 658\text{nm}.$$

19.21 Another Neodymium-YAG laser

a. Each photon has an energy given by

$$E = \frac{hc}{\lambda} = \frac{6.63 \times 10^{-34}\text{Js} \times 3 \times 10^8 \frac{\text{m}}{\text{s}}}{532 \times 10^{-9}\text{m}} = 3.74 \times 10^{-19}\text{J per photon. Similarly the}$$

momentum is $p = \frac{h}{\lambda} = \frac{6.63 \times 10^{-34}\text{Js}}{532 \times 10^{-9}\text{m}} = 1.25 \times 10^{-27} \frac{\text{kgm}}{\text{s}}$ per photon.

b. Since the total energy is 5J, we calculate the number of photons by dividing the total energy by the energy per photon. Thus we have

$$\# = \frac{E_{\text{total}}}{E/\text{photon}} = \frac{5\text{J}}{3.74 \times 10^{-19}\text{J/photon}} = 1.34 \times 10^{19}\text{ photons.}$$

c. In 1 ns, if all the photons in part b are absorbed, each carrying a momentum p , then the force is the total change in momentum and is given by

$$F = N \frac{\Delta p}{\Delta t} = 1.34 \times 10^{19} \times \frac{1.25 \times 10^{-27} \frac{\text{kgm}}{\text{s}}}{1 \times 10^{-9}\text{s}} = 16.7\text{N}.$$

24.7 For a $1.2 \times 10^6\text{eV} \times \frac{1.6 \times 10^{-19}\text{J}}{1\text{eV}} = 1.92 \times 10^{-13}\text{J photon}$,

a. its momentum is given by $p = \frac{E}{c} = \frac{1.92 \times 10^{-13}\text{J}}{3 \times 10^8 \frac{\text{m}}{\text{s}}} = 6.4 \times 10^{-22} \frac{\text{kgm}}{\text{s}}$.

b. its wavelength is given by the de Broglie relation

$$\lambda = \frac{h}{p} = \frac{6.63 \times 10^{-34}\text{Js}}{6.4 \times 10^{-22} \frac{\text{kgm}}{\text{s}}} = 1.04 \times 10^{-12}\text{m}.$$

c. its frequency is given by $f = \frac{c}{\lambda} = \frac{3 \times 10^8 \frac{\text{m}}{\text{s}}}{1.04 \times 10^{-12}\text{m}} = 2.9 \times 10^{20}\text{s}^{-1}$.

24.1 Relativistic energy and momentum for an object of mass m .

For an object with a $m = 1\text{kg}$ rest mass, it has a rest energy of $E = mc^2 = 9 \times 10^{16}\text{J}$.

The Lorentz factor is given by: $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$, with the relativistic momentum $p =$

γmv , and the relativistic energy $E^2 = p^2 c^2 + m^2 c^4$.

- a. For a velocity of $0.8c$, $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{\sqrt{1 - 0.64}} = 1.67$. The relativistic

momentum and energy are therefore,

$$p = \gamma mv = 1.67 \times 1 \text{ kg} \times 0.8 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 4.0 \times 10^8 \frac{\text{kgm}}{\text{s}} \text{ and}$$

$$E = \sqrt{p^2 c^2 + m^2 c^4} = \sqrt{(4.0 \times 10^8 \frac{\text{kgm}}{\text{s}} \times 3 \times 10^8 \frac{\text{m}}{\text{s}})^2 + (9 \times 10^{16} \text{ J})^2} = 1.5 \times 10^{17} \text{ J}.$$

- b. Following the procedure in part a, for a velocity of $0.9c$,

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{\sqrt{1 - 0.81}} = 2.29, \text{ the relativistic momentum is } 6.19 \times 10^8 \text{ kgm/s,}$$

and the relativistic energy is $2.07 \times 10^{17} \text{ J}$.

- c. For a velocity of $0.95c$, $\gamma = 3.20$, the relativistic momentum is $9.13 \times 10^8 \text{ kgm/s}$, and the relativistic energy is $2.88 \times 10^{17} \text{ J}$.
- d. For a velocity of $0.99c$, $\gamma = 7.09$, the relativistic momentum is $2.11 \times 10^9 \text{ kgm/s}$, and the relativistic energy is $6.387 \times 10^{17} \text{ J}$.
- e. For a velocity of $0.999c$, $\gamma = 22.4$, the relativistic momentum is $6.70 \times 10^9 \text{ kgm/s}$, and the relativistic energy is $2.0 \times 10^{18} \text{ J}$.

- 24.2 For an electron with rest mass $9.11 \times 10^{-31} \text{ kg}$, it has a rest energy of $E = mc^2 = 8.199 \times 10^{-14} \text{ J} = 0.511 \text{ MeV}$. The Lorentz factor is given by: $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$, and

the relativistic momentum and energy are given respectively as $p = \gamma mv$ and

$$E = \sqrt{p^2 c^2 + m^2 c^4}.$$

- a. For the electron with a velocity of $0.8c$, $\gamma = 1.67$. The relativistic momentum and energy are therefore,

$$p = \gamma mv = 1.67 \times 9.11 \times 10^{-31} \text{ kg} \times 0.8 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 3.65 \times 10^{-22} \frac{\text{kgm}}{\text{s}} \text{ and}$$

$$E = \sqrt{p^2 c^2 + m^2 c^4} = \sqrt{(3.65 \times 10^{-22} \frac{\text{kgm}}{\text{s}} \times 3 \times 10^8 \frac{\text{m}}{\text{s}})^2 + (8.199 \times 10^{-14} \text{ J})^2} = 1.37 \times 10^{-13} \text{ J}$$

Using the fact that $1.6 \times 10^{-19} \text{ J} = 1 \text{ eV}$, the relativistic energy is given as

$$E = 1.37 \times 10^{-13} \text{ J} \times \frac{1 \text{ eV}}{1.6 \times 10^{-19} \text{ J}} = 0.855 \times 10^6 \text{ eV} = 0.855 \text{ MeV}. \text{ Further it can be}$$

shown that the total relativistic energy is given as $E = \gamma mc^2 = \gamma E_{rest}$. Thus the relativistic energy can be computed in a more efficient method.

$$E = \gamma E_{rest} = 1.67 \times 0.511 \text{ MeV} = 0.855 \text{ MeV}.$$

- b. Following the method outlined in part a, for a velocity of $0.9c$, $\gamma = 2.29$. The relativistic momentum is therefore,

$$p = \gamma mv = 2.29 \times 9.11 \times 10^{-31} \text{ kg} \times 0.9 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 5.63 \times 10^{-22} \frac{\text{kgm}}{\text{s}} \text{ and}$$

$$E = \gamma E_{rest} = 2.29 \times 0.511 \text{ MeV} = 1.17 \text{ MeV}.$$

- c. For a velocity of $0.95c$, $\gamma = 3.2$. The relativistic momentum is therefore,
 $p = \gamma mv = 3.2 \times 9.11 \times 10^{-31} \text{ kg} \times 0.95 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 8.31 \times 10^{-22} \frac{\text{kgm}}{\text{s}}$ and
 $E = \gamma E_{rest} = 3.2 \times 0.511 \text{ MeV} = 1.63 \text{ MeV}$.
- d. For a velocity of $0.99c$, $\gamma = 7.09$. The relativistic momentum is therefore,
 $p = \gamma mv = 7.09 \times 9.11 \times 10^{-31} \text{ kg} \times 0.99 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 1.92 \times 10^{-21} \frac{\text{kgm}}{\text{s}}$ and
 $E = \gamma E_{rest} = 7.09 \times 0.511 \text{ MeV} = 3.62 \text{ MeV}$.
- e. For a velocity of $0.999c$, $\gamma = 22.4$. The relativistic momentum is therefore,
 $p = \gamma mv = 22.4 \times 9.11 \times 10^{-31} \text{ kg} \times 0.999 \times 3 \times 10^8 \frac{\text{m}}{\text{s}} = 6.12 \times 10^{-21} \frac{\text{kgm}}{\text{s}}$ and
 $E = \gamma E_{rest} = 22.4 \times 0.511 \text{ MeV} = 11.45 \text{ MeV}$.

24.4 The relativistic momentum and relativistic energy are given as $p = \gamma mv$ and $E = \gamma mc^2$ respectively. To show the relation between energy and momentum, equation (24.8), we start by squaring the relativistic energy. This gives us $E^2 = \gamma^2 m^2 c^4$. Next, we use a mathematical “trick.” We add and subtract the same quantity from the right hand side of the equation we just developed. The quantity we want to add and subtract is v^2 . This produces factoring out a factor of c^2 , $E^2 = \gamma^2 m^2 c^2 (c^2 + v^2 - v^2)$. Expanding this result we get $E^2 = \gamma^2 m^2 c^2 v^2 + \gamma^2 m^2 c^2 (c^2 - v^2)$. Recognizing that the first term is nothing more than $p^2 c^2$ allows us to write $E^2 = p^2 c^2 + \gamma^2 m^2 c^2 (c^2 - v^2)$. Factoring out a c^2 from the 2nd term on the right hand side give us $E^2 = p^2 c^2 + \gamma^2 m^2 c^4 \left(1 - \frac{v^2}{c^2}\right)$.

The quantity $\left(1 - \frac{v^2}{c^2}\right)$ is simply $\frac{1}{\gamma^2}$. Therefore we arrive at the desired result,
 $E^2 = p^2 c^2 + m^2 c^4$.